

REMARKS

Claim 1 remains in the application. Claim 1 has been amended. Claims 19, 20, 26, and 27 have been canceled. Claim 30 has been added. The subject matter of claims 26 and 27 will be re-inserted in a divisional application containing original claims 21 and 22. Reconsideration of this application, as amended, is respectfully requested.

Claim 1 has been amended to include the phrase "at least one" so that the final clause conforms to clauses a), c), and d). Claim 1 has been further amended to specify that all of the steps of the method are carried out in aqueous medium. Support for this amendment can be found at page 5, paragraph 0023 of the specification. Claim 30 has been added to specify that the molecular weight of the First Macromolecule is at least 2,000 daltons. The remainder of the subject matter of claim 30 is identical to the subject matter of claim 1, as filed in the AMENDMENT AND RESPONSE filed December 16, 2004. Support for this amendment can be found at page 16, paragraph 0055 of the specification and in paragraphs 0037, 0049, 0050, 0051, 0052, 0056, 0060 of the specification, as described in the AMENDMENT AND RESPONSE filed December 16, 2004 .

Claim 1 stands rejected under 35 U. S. C. §102(e) as being anticipated by Yu et al. (U. S. 6,492,105). This rejection is respectfully traversed for the following reasons.

Yu et al., U. S. Patent No. 6,492,105 (hereinafter "Yu et al."), discloses polypeptide binding molecules prepared using solid phase peptide synthesis. Solid-phase synthesis begins at the carboxy-terminus of the putative polypeptide by coupling a protected amino acid to a suitable resin, which reacts with the carboxy group of the C-terminal amino acid to form a bond that is readily cleaved later, such as a halomethyl resin, hydroxymethyl resin, aminomethyl resin, benzhydrylamine resin, or t-alkyloxycarbonyl-hydrazide resin. After removal of the α -amino protecting group with, for example, trifluoroacetic acid in methylene chloride and neutralizing in, for example, TEA, the next cycle in synthesis is ready to proceed. The remaining α -amino and, if necessary, side-chain-protected amino acids are then coupled

sequentially in the desired order by condensation to obtain an intermediate compound connected to the resin. After the desired amino acid sequence has been completed, the intermediate polypeptide is removed from the resin support by treatment with a reagent, such as liquid HF and one or more thio-containing scavengers, which not only cleaves the polypeptide from the resin, but also cleaves all the remaining side-chain protecting groups. Following HF cleavage, the protein sequence is washed with ether, transferred to a large volume of dilute acetic acid, and stirred at pH adjusted to about 8.0 with ammonium hydroxide. Upon pH adjustment, the polypeptide takes its desired conformational arrangement.

Although Yu et al., at column 12, lines 61-63, discloses the presence of a covalent bond between an amino acid (which the Examiner has deemed analogous to the First Macromolecule recited in claim 1 of the present application) and another amino acid (which the Examiner has deemed analogous to the Second Macromolecule recited in claim 1 of the present application), Yu et al. is merely describing a conventional variant of synthesizing a polypeptide in which a preformed oligopeptide is coupled to a growing solid phase polypeptide chain in order to produce a longer, but still conventional, polypeptide chain. The synthesis described in Yu et al. is carried out under conditions that are substantially different from those conditions employed in the method of the present invention, with the result that the composition produced by the method described in Yu et al. is substantially different from the composition produced by the method of the present invention.

According to Yu et al., hydrazide-containing peptides were immobilized by facilitating hydrazone bond formation, the Gl-1 and -2 peptides were immobilized via reductive amination using NaCNBH_3 . The nine peptides were evaluated by affinity chromatography for their ability to capture the REFACTO[®] described in Example I, under specific binding and release conditions. See column 18, line 67 through column 19, line 3 and column 19, lines 9-11 of Yu et al. REFACTO[®] is a recombinantly produced factor VIII-like polypeptide consisting of two segments of human factor VIII. See column 15, lines 3-10 of Yu et al.

The hydrazide-containing peptide is considered to be the first macromolecule, while the recombinantly produced factor VIII-like polypeptide consisting of two segments of human factor VIII for which the hydrazide-containing peptide has binding affinity is considered to be the second macromolecule. The method described in Yu et al. uses this binding affinity to isolate the factor VIII-like polypeptide from a complex mixture by binding the factor VIII-like polypeptide to a hydrazide-containing peptide immobilized on a solid support, and then releasing the factor VIII-like polypeptide by disruption of the bond between the factor VIII-like polypeptide and hydrazide-containing peptide. Yu et al. does not disclose or suggest that the bond to be disrupted is the bond that links the hydrazide-containing peptide to the solid support. In fact, the release conditions are specifically chosen in order to dissociate the hydrazide-containing polypeptide bound to the solid support from the factor VIII-like polypeptide, as pointed out in column 8, lines 15-44 of Yu et al. Further, in Example III, Yu et al. demonstrates that the solid support linked via hydrazone groups to several polypeptides shows binding and release of the factor VIII-like polypeptide. Yu et al. then shows that the solid support can be successfully reused without further exposure to the hydrazide-containing peptide, thereby showing that the peptide-hydrazone linkage to the solid support remains intact through the binding, elution, and washing steps described in Yu et al.

In contrast, in the method of the present invention, the linkage targeted for disruption is the linkage between the solid support and the complex comprising the First Macromolecule and the at least one Second Macromolecule. For this reason, it is submitted that Yu et al. does anticipate claim 1 of the present application.

Claims 19 and 20 stand rejected under 35 U. S. C. §103(a) as being unpatentable over Yu et al. (U. S. 6,492,105). Claims 19 and 20 have been canceled.

Claims 1, 19, and 20 stand rejected under 35 U. S. C. §102(e) as being anticipated by Schwartz (U. S. 2003/0013857 A1). This rejection is respectfully traversed for the following reasons.

Schwartz, U.S. Patent Application Publication 2003/0013857 A1 (hereinafter "Schwartz"), discloses modified solid supports that include solid

supports that have been modified by reaction with a bifunctional reagent that possess a hydrazine or oxyamino group. These modified solid supports are useful in immobilization of biomolecules that possess or are modified to possess a carbonyl group. In one embodiment, aliphatic bifunctional hydrazide reagents are provided. These reagents include a cleavable bond for further manipulation. Cleavable bonds include, but are not limited to, acid cleavable, photocleavable and disulfide bonds.

In Schwartz, paragraph 0110 describes the use of cleavable linkers to create a drug-antibody conjugate, which is cleaved by physiological processes following endocytosis. Schwartz also refers to the use of cleavable disulfide linkages to isolate receptors following covalent linking between a ligand and a receptor. Paragraph 0111 of Schwartz describes the use of the bifunctional hydrazides to modify biomolecules or carriers in a single step. These modified aliphatic hydrazide molecules or carriers can be subsequently reacted with carbonyl containing biomolecules, drug, or other therapeutic or diagnostic reagent to form a hydrazone that can be cleaved following exposure to mild aqueous acid conditions. Paragraph 0112 of Schwartz describes that solid supports such as beads, chromatographic supports or surfaces are modified with aliphatic hydrazide reagents. Paragraph 0118 of Schwartz describe solid supports modified with the reagents described therein so as to allow immobilization of biomolecules that contain a hydrazide group. Paragraph 0147 of Schwartz describes hydrazino modified beads for forming stable hydrazones when reacted with molecules possessing carbonyl groups. Paragraph 0172 of Schwartz describes hydrazones that are prepared from bifunctional hydrazine intermediates by treatment of these compounds with an aliphatic aldehyde or ketone to protect the hydrazine as its hydrazone. Paragraph 0175 of Schwartz describes conjugates formed by the reaction of hydrazine-modified molecules with carbonyl-containing or carbonyl-modified molecules, hydrazine-modified molecules with carbonyl-modified surfaces or carbonyl-containing or carbonyl-modified molecules with hydrazine-modified surfaces. Paragraph 0176 of Schwartz describes the modification of an organic soluble natural or synthetic molecule such as a hydrophobic peptide in organic solution by direct addition of the protected bifunctional hydrazino molecule to an organic soluble compound possessing an amine or thiol

substituted moiety. Paragraph 0177 of Schwartz describes the use of the reagents described therein to form crosslinks between a wide variety of molecules, such as, for example, protein-protein conjugates or protein-polymer conjugates. Paragraph 0178 of Schwartz describes the crosslinking of conjugated biomolecules by incubation at ambient temperature of a hydrazine-modified biomolecule and a carbonyl-modified biomolecule. Paragraph 0179 of Schwartz describes immobilization of biomolecules to surfaces using a crosslinking couple by modifying the biomolecule with either a hydrazino, oxyamino, or a carbonyl moiety and contacting the modified biomolecule to a surface possessing its reactive partner, e.g., a hydrazino or oxyamino moiety for a carbonyl-modified biomolecule, or a carbonyl moiety for a hydrazino- or oxyamino-modified biomolecule. Paragraph 0180 of Schwartz describes oxidation with sodium periodate of the carbohydrate domain of an antibody that is not involved in antigen binding to yield aldehyde moieties. The aldehyde moieties react directly with hydrazine- or oxyamino-modified surfaces to yield a stable covalent linkage in which a large percentage of the antibody active sites are available for antigen binding. Paragraph 0181 of Schwartz describes that bisarylhydrazones formed from the reaction of an aromatic hydrazine and an aromatic carbonyl exhibit more stability than do hydrazones formed from the reaction of aliphatic hydrazines and aliphatic carbonyls. Example 17 of Schwartz exemplifies the coupling of hydrazine-modified BSA to oxidized dextran.

Although Schwartz provides a great number of descriptions of conjugations between biomolecules and solid supports with cleavable linkers, Schwartz does not disclose or suggest a method for preparing a conjugate in which (1) a first macromolecule is linked to a solid surface through a cleavable linker, (2) a second macromolecule is linked to the first macromolecule through a non-cleavable linker, and (3) the linker between the solid surface and the first macromolecule is cleaved, thereby (4) releasing the conjugate consisting of the first macromolecule linked to the second macromolecule by means of a non-cleavable linker. For these reasons, Schwartz does not anticipate claim 1.

Claims 19 and 20 have been canceled.

Claim 1 stands rejected under 35 U. S. C. §103(a) as being unpatentable over Merrifield (Reference AK of PTO form 1449). This rejection is respectfully traversed for the following reasons.

Merrifield, Solid Phase Peptide Synthesis. I. The Synthesis of a Tetrapeptide, Journal of American Chemical Society, USA, 85, pp. 2149-2154 (July 1963) (hereinafter "Merrifield"), discloses the stepwise addition of protected amino acids to a growing peptide chain which was bound by a covalent bond to a solid resin particle. This provided a procedure whereby reagents and by-products were removed by filtration, and the recrystallization of intermediates was eliminated.

The method described in Merrifield depends on the attachment of the first amino acid of the chain to a solid polymer by a covalent bond, the addition of the succeeding amino acids one at a time in a stepwise manner until the desired sequence is assembled, and finally the removal of the peptide from the solid support. The intermediate peptides are purified, not by recrystallization procedures, but by dissolving away the impurities.

In the present invention, a First Macromolecule is attached to a Second Macromolecule, and so forth until a chain of macromolecules is formed. Because individual amino acids are not macromolecules, it is submitted that Merrifield does not render claim 1 obvious to one of ordinary skill in the art.

The method described in Merrifield differs from the method claimed in the present application in the following respects:

- (1) the nature of the materials being coupled;
- (2) the nature of the chemical bonds being formed and broken;
- (3) the nature of the solvents used;
- (4) the reagents used in the reactions;
- (5) the reaction conditions; and
- (6) the character of the products produced

In Merrifield, the reactions are performed under conditions typical of organic synthesis, that is, with solvents such as, for example, dimethylformamide,

acetic acid, ethanol, and triethylamine. See page 2152, column 2 of Merrifield. Claim 1 has been amended to specify that each step of the method is performed under aqueous conditions. To retain their biological activities, the macromolecules used in the present invention must remain in aqueous solvents, preferably under substantially physiological conditions. The organic synthetic conditions required for effective synthesis by means of the process described in Merrifield would denature most proteins, thereby destroying their biological activities and rendering the products useless for the purpose intended.

In Merrifield, the amino acid derivatives that are being coupled are not macromolecules. In the method of the present invention, the materials being coupled are macromolecules, which are much larger than the amino acid derivatives used in synthesis of peptides. Accordingly, claim 30 has been added to specify this feature. In view of the changes to claim 1 and the features of new claim 30, it is submitted that Merrifield does not render claims 1 and 30 obvious to one of ordinary skill in the art.


Claims 19, 20, 26, and 27 were rejected under 35 U. S. C. § 102/103 as being anticipated by obvious over each of Xu et al (US 6,737,240), Rice et al (US 6,759,036), or Cook et al (US 6,753,423). Claims 19, 20, 26, and 27 have been canceled.

In view of the foregoing, it is submitted that claim 1, as amended, and new claim 30 are in condition for allowance, and official Notice of Allowance is respectfully requested.

23492

Abbott Laboratories
D-377 AP6A-1
100 Abbott Park Road
Abbott Park, Illinois 60064-3500
Telephone: (847) 937-6182

Respectfully submitted,
John C. Russell


David L. Weinstein
Registration No. 28, 128
Attorney for Applicants